

Enumerations for Permutations by Circular Peak Sets

Pierre Bouchard^a Hungyung Chang^b Jun Ma^{c,*} Jean Yeh^{d,†}

June 5, 2008

^a Dépt. de mathématiques, Université du Québec à Montréal, C.P. 8888, Succ. Centre-Ville Montréal,
Canada H3C 3P8

^{b,c} Institute of Mathematics, Academia Sinica, Taipei, Taiwan

^d Department of Mathematics, National Taiwan University, Taipei, Taiwan

Abstract

The circular peak set of a permutation σ is the set $\{\sigma(i) \mid \sigma(i-1) < \sigma(i) > \sigma(i+1)\}$.

In this paper, we focus on the enumeration problems for permutations by circular peak sets. Let $cp_n(S)$ denote the number of the permutations of order n which have the circular peak set S . For the case with $|S| = 0, 1, 2$, we derive the explicit formulas for $cp_n(S)$. We also obtain some recurrence relations for the sequence $cp_n(S)$ and give the formula for $cp_n(S)$ in the general case.

Keywords: Circular peak; Lattice path; Permutation;

*Email address of the corresponding author: majun@math.sinica.edu.tw

†jean.yh@ms45.url.com.tw

1 Introduction

Throughout this paper, let $[m, n] := \{m, m+1, \dots, n\}$, $[n] := [1, n]$ and $[m, n] = \emptyset$ if $m > n$. Let $\mathfrak{S}_n$ be the set of all the permutations in the set $[n]$. We will write permutations of $\mathfrak{S}_n$ in the form $\sigma = (\sigma(1)\sigma(2)\cdots\sigma(n))$. We say that a permutation σ has a circular descent of value $\sigma(i)$ if $\sigma(i) > \sigma(i+1)$ for any $i \in [n-1]$. The circular descent set of a permutation σ , denoted $CDES(\sigma)$, is the set $\{\sigma(i) \mid \sigma(i) > \sigma(i+1)\}$. For any $S \subseteq [n]$, define a set $CDES_n(S)$ as $CDES_n(S) = \{\sigma \in \mathfrak{S}_n \mid CDES(\sigma) = S\}$ and use $cdes_n(S)$ to denote the number of the permutations in the set $CDES_n(S)$, i.e., $cdes_n(S) = |CDES_n(S)|$. In a join work [2], Hungyung Zhang et al. derive the explicit formula for $cdes_n(S)$. As a application of the main results in [2], they also give the enumeration of permutation tableaux according to their shape and generalizes the results in [4]. Moreover, Robert J. Clarke et al. [3] gave the conceptions of linear peak and cyclic peak and studied some new Mahonian permutation statistics. In this paper, we say that a permutation σ has a *circular peak* of value $\sigma(i)$ if $\sigma(i-1) < \sigma(i) > \sigma(i+1)$ for any $i \in [2, n-1]$. The *circular peak set* of a permutation σ , denoted $CP(\sigma)$, is the set $\{\sigma(i) \mid \sigma(i-1) < \sigma(i) > \sigma(i+1)\}$. For example, the circular peak set of $\sigma = (48362517)$ is $\{5, 6, 8\}$. Since σ has no circular peaks when $n \leq 2$, we always suppose that $n \geq 3$. For any $S \subseteq [n]$, we define a set $CP_n(S)$ as $CP_n(S) = \{\sigma \in \mathfrak{S}_n \mid CP(\sigma) = S\}$. Obviously, if $\{1, 2\} \subseteq S$, then $CP_n(S) = \emptyset$.

Example 1.1

$$CP_5(\{4, 5\}) = \{ \quad 14253, 14352, 24153, 34152, 24351, 34251, \\ 15243, 15342, 25143, 35142, 25341, 35241 \quad \}$$

Suppose that $S = \{i_1, i_2, \dots, i_k\}$, where $i_1 < i_2 < \dots < i_k$. Pierre Bouchard et al. [1] proved that the necessary and sufficient conditions for $CP_n(S) \neq \emptyset$ are $i_j \geq 2j + 1$ for all $j \in [k]$.

Let $\mathcal{P}_n = \{S \mid CP_n(S) \neq \emptyset\}$. We can make the set $\mathcal{P}_n$ into a poset $\mathcal{P}_n$ by defining $S \preceq T$ if $S \subseteq T$ as sets. See [1] for the properties of the poset $\mathcal{P}_n$.

In this paper, we focus on the enumerations for permutations by circular peak sets. Let $cp_n(S)$ denote the number of the permutations in the sets $CP_n(S)$, i.e., $cp_n(S) = |CP_n(S)|$. For the case with $|S| = 0, 1, 2$, we derive the explicit formulas for $cp_n(S)$. For the general case, we consider the recurrence relations for $cp_n(S)$. We find that if $n \geq 3$ and $S \subseteq [n - 1]$ with $CP_n(S) \neq \emptyset$, then $cp_n(S)$ satisfies the following recurrence relation

$$cp_n(S \cup \{n\}) = [n - 2 - 2|S|]cp_{n-1}(S) + \sum_{j \notin S} 2cp_{n-1}(S \cup \{j\}).$$

But we are more interested in another recurrence relation as follows:

suppose $n \geq 3$, $k \geq 0$ and $S \subseteq [n - k - 1]$ with $CP_n(S) \neq \emptyset$, then

$$cp_n(S \cup [n - k + 1, n]) = 2(k + 1)cp_{n-1}(S \cup [n - k, n - 1]) + k(k + 1)cp_{n-2}(S \cup [n - k, n - 2]).$$

First, let $k \geq 1$, $n \geq \max\{3, 2k\}$ and $CP_n([n - k + 1, n]) \neq \emptyset$, by this recurrence relation, we conclude that

$$cp_n([n - k + 1, n]) = k(k + 1) \sum_{i=1}^k (-1)^{i+1} b_{k,i} [(2k + 2)^{n-2k} - (2k + 2 - 2i)^{n-2k}],$$

where the coefficients $b_{k,i}$ satisfy the following recurrence relation:

$$b_{k+1,i} = \begin{cases} k(k + 1)^2 \sum_{j=1}^k (-1)^{j+1} b_{k,j} & \text{if } i = 1 \\ k(k + 1) \frac{k+2-i}{i} b_{k,i-1} & \text{if } 2 \leq i \leq k + 1 \end{cases}$$

with initial condition $b_{1,1} = \frac{1}{2}$. And then we study the connections between this recurrence relation and the circular-peak path with weight. A *circular-peak path* is a lattice path in the first quadrant starting at $(r, 0)$ and ending at (n, k) with only two kinds of steps—*horizon step* $H = (1, 0)$ and *rise step* $R = (2, 1)$. We consider a circular-peak path P from $(r, 0)$ to (n, k) as a word of $n - r - k$ letters using only H and R . Let $P_{r,n,k}$ be the set of all the circular-peak paths from the vertices $(r, 0)$ to (n, k) . Given an integer i and $P = e_1 e_2 \cdots e_{n-k-r} \in P_{r,n,k}$, if the step e_j connects the vertices (x, y) with $(x + 1, y)$, then the weight of e_j , denoted $w_i(e_j)$, is $2i + 2(y + 1)$; if the step e_j connects the vertices (x, y) with $(x + 2, y + 1)$, then the weight of e_j , denoted $w_i(e_j)$, is $(y + i + 1)(y + i + 2)$; at last, let $w_i(P) = \prod_{j=1}^{n-k-r} w_i(e_j)$ be the weight of the circular-peak path P and $w(i, r, n, k) = \sum_{P \in P_{r,n,k}} w_i(P)$. It is proved that if $k \geq 0$, $n \geq k + 4$, $S \subseteq [3, n - k - 1]$ with $CP_n(S) \neq \emptyset$ and $r = \max S$, then

$$cp_n(S \cup [n - k + 1, n]) = \sum_{i=0}^k w(i, r, n - i, k - i) cp_{r+i}(S \cup [r + 1, r + i]).$$

We must give the formula for $w(i, r, n, k)$ as follows:

$$w(i, r, n, k) = 2^{n-r-2k} \prod_{m=0}^{k-1} (m + i + 1)(m + i + 2) \sum \prod_{m=0}^k [i + m + 1]^{t_m},$$

where the sum is over all $(k+1)$ -tuples $(t_0, t_1, \dots, t_k)$ such that $\sum_{m=0}^k t_m = n - r - 2k$ and $t_m \geq 0$.

For any $S \subseteq [3, n]$, define the *type* of the set S as $(r_1^{k_1}, r_2^{k_2}, \dots, r_m^{k_m})$ if $S = \bigcup_{i=1}^m [r_i - k_i + 1, r_i]$

such that $r_i \leq r_{i+1} - k_{i+1} - 1$ for all $i \in [1, m - 1]$. We may state one of the main results of

the paper as follows:

$$\begin{aligned} cp_n(S) &= 2^{n-r_m} \sum_{i_1=0}^{k_m} \sum_{i_2=0}^{k_{m-1}+i_1} \cdots \sum_{i_{m-1}=0}^{k_2+i_{m-2}} \prod_{j=1}^{m-1} \\ &\quad w(i_j, r_{m-j}, r_{m-j+1} + i_{j-1} - i_j, k_{m-j+1} + i_{j-1} - i_j) cp_{r_1+i_{m-1}}([r_1 - k_1 + 1, r_1 + i_{m-1}]), \end{aligned}$$

where $i_0 = 0$, when $m \geq 2$ and $CP_n(S) \neq \emptyset$.

The paper is organized as follows. In Section 2, we will consider the enumerations of the permutations in the sets $CP_n(S)$ with $|S| = 0, 1, 2$. In Section 3, we will derive some recurrence relations for the sequences $cp_n(S)$ and give the formula for $cp_n(S)$ in the general case. In the Appendix, we list all the values of $cp_n(S) > 0$ for $3 \leq n \leq 8$.

2 The Enumerations for The Permutations in The Set

$CP_n(S)$ with $|S| = 0, 1, 2$

In this section, we will consider the enumeration problems of the permutations in the sets $CP_n(S)$ with $|S| = 0, 1, 2$.

Let $cp_n(S)$ denote the number of the elements in the sets $CP_n(\{S\})$, i.e., $cp_n(S) = |CP_n(S)|$. First, we need the following lemma.

Lemma 2.1 *Suppose $n \geq 3$ and $S \subseteq [n]$ with $CP_n(S) \neq \emptyset$. Then*

- (1) $cp_{n+1}(S) = 2cp_n(S)$, and
- (2) let $m = \max S$, then $cp_n(S) = 2^{n-m}cp_m(S)$ for any $n \geq m$.

Proof. (1) It is easy to check that $((n+1)\sigma(1) \cdots \sigma(n)) \in CP_{n+1}(S)$ and $(\sigma(1) \cdots \sigma(n)(n+1)) \in CP_{n+1}(S)$ for any $\sigma = (\sigma(1) \cdots \sigma(n)) \in CP_n(S)$. Conversely, for any $\sigma \in CP_{n+1}(S)$, the position of the letter $n+1$ is 1 or $n+1$, i.e., $\sigma^{-1}(n+1) = 1$ or $n+1$, since $n+1 \notin S$. Hence, $cp_{n+1}(S) = 2cp_n(S)$.

- (2) Iterating the identity in (1), we immediately obtain that $cp_n(S) = 2^{n-m}cp_m(S)$. ■

For any $\sigma \in \mathfrak{S}_n$, let τ be a subsequence $(\sigma(j_1)\sigma(j_2)\cdots\sigma(j_k))$ of $(\sigma(1)\cdots\sigma(n))$, where $1 \leq j_1 < j_2 < \cdots < j_k \leq n$. Define $red_{\sigma,\tau}$ as an increasing bijection of $\{\sigma(j_1), \sigma(j_2), \dots, \sigma(j_k)\}$ onto $[k]$. Let $red_{\sigma}(\tau) = (red_{\sigma,\tau}(\sigma(j_1))red_{\sigma,\tau}(\sigma(j_2))\cdots red_{\sigma,\tau}(\sigma(j_k)))$.

Theorem 2.1 *Let $n \geq 3$. Then (1) $cp_n(\emptyset) = 2^{n-1}$,*

(2) $cp_n(\{i\}) = 2^{n-2}(2^{i-2} - 1)$ for any $i \in [3, n]$, and

(3) $cp_n(\{i, j\}) = 2^{n-3}(2^{i-2} - 1)(2^{j-i-1} - 1) + 2^{n+j-i-5} \cdot 3(3^{i-2} - 2^{i-1} + 1)$ for any $i, j \in [3, n]$

and $i < j$.

Proof. (1) For any $\sigma \in \mathfrak{S}_n$, suppose that the position of the letter 1 is $i + 1$, i.e., $\sigma^{-1}(1) = i + 1$. Then $\sigma \in CP_n(\emptyset)$ if and only if σ has the form $\sigma(1) > \cdots > \sigma(i + 1) < \cdots < \sigma(n)$. For each letter $j \neq 1$, the position of j has two possibilities at the left or right of 1. Hence, $cp_n(\emptyset) = 2^{n-1}$.

(2) By Lemma 2.1(2), we first consider the number of the permutations in the set $CP_i(\{i\})$, where $i \geq 3$. For any $\sigma \in CP_i(\{i\})$, suppose that the position of the letter i is $k + 1$, i.e., $\sigma^{-1}(i) = k + 1$, then $1 \leq k \leq i - 2$, $red_{\sigma}(\sigma(1)\cdots\sigma(k)) \in CP_k(\emptyset)$ and $red_{\sigma}(\sigma(k + 2)\cdots\sigma(i)) \in CP_{i-k-1}(\emptyset)$. There are $\binom{i-1}{k}$ ways to form the set $\{\sigma(1), \dots, \sigma(k)\}$. So, $cp_i(\{i\}) = \sum_{k=1}^{i-2} \binom{i-1}{k} 2^{k-1} 2^{i-k-2} = 2^{i-2}(2^{i-2} - 1)$. Hence, $cp_n(\{i\}) = 2^{n-2}(2^{i-2} - 1)$.

(3) It is easy to check that the identity holds when $i = 3$ and $j = 4$. By Lemma 2.1(2), we first consider the number of the permutations in the set $CP_j(\{i, j\})$, where $3 \leq i < j$.

We begin from the case $\sigma \in CP_j(\{i, j\})$ with $\sigma^{-1}(i) < \sigma^{-1}(j)$. Let

$$T_1(\sigma) = \{\sigma(k) \mid \sigma(k) < i \text{ and } k < \sigma^{-1}(i)\},$$

$$T_2(\sigma) = \{\sigma(k) \mid \sigma(k) < i \text{ and } \sigma^{-1}(i) < k < \sigma^{-1}(j)\},$$

$$T_3(\sigma) = \{\sigma(k) \mid \sigma(k) < i \text{ and } k > \sigma^{-1}(j)\}.$$

Note that $T_k(\sigma) \neq \emptyset$ for $k = 1, 2$ since σ must have a circular peak i and $\bigcup_{k=1}^3 T_k(\sigma) = [i - 1]$.

Let

$$T_4(\sigma) = \{\sigma(k) \mid i < \sigma(k) < j \text{ and } k < \sigma^{-1}(i)\}$$

$$T_5(\sigma) = \{\sigma(k) \mid i < \sigma(k) < j \text{ and } \sigma^{-1}(i) < k < \sigma^{-1}(j)\}.$$

We discuss the following two subcases.

Subcase 1. $T_3(\sigma) = \emptyset$

Let $T_6(\sigma) = \{\sigma(k) \mid i < \sigma(k) < j, k > \sigma^{-1}(j)\}$. Then $T_6(\sigma) \neq \emptyset$ since σ must have a circular peak j and $\bigcup_{k=4}^6 T_k(\sigma) = [i + 1, j - 1]$. For $k = 1, 2, 6$, the subsequences of σ , which is determined by the elements in $T_k(\sigma)$, corresponds to a permutation in $CP_{|T_k(\sigma)|}(\emptyset)$. The subsequences of σ , which is determined by the elements in $T_4(\sigma)$ and $T_5(\sigma)$, are decreasing and increasing, respectively. So, the number of the permutations under this subcase is

$$\sum_{(T_1, T_2)} \binom{i-1}{|T_1|, |T_2|} 2^{|T_1|-1} 2^{|T_2|-1} \sum_{(T_4, T_5, T_6)} \binom{j-i-1}{|T_4|, |T_5|, |T_6|} 2^{|T_6|-1} = 2^{j-4} (2^{i-2} - 1) (2^{j-i-1} - 1),$$

where the first sum is over all pairs (T_1, T_2) such that $T_i \neq \emptyset$ for $i = 1, 2$ and $T_1 \cup T_2 = [i - 1]$;

the second sum is over all triples (T_4, T_5, T_6) such that $T_6 \neq \emptyset$ and $T_4 \cup T_5 \cup T_6 = [i + 1, j - 1]$.

Subcase 2. $T_3(\sigma) \neq \emptyset$

Suppose that $\min T_3(\sigma) = s$, let

$$T_6(\sigma) = \{\sigma(k) \mid i < \sigma(k) < j \text{ and } \sigma^{-1}(j) < k < \sigma^{-1}(s)\}$$

$$T_7(\sigma) = \{\sigma(k) \mid i < \sigma(k) < j \text{ and } k > \sigma^{-1}(s)\}.$$

Then, for $k = 1, 2, 3$, the subsequences of σ , which is determined by the elements in $T_k(\sigma)$, corresponds to a permutation in $CP_{|T_k(\sigma)|}(\emptyset)$. The subsequences of σ , which is determined by the elements in $T_4(\sigma)$ and $T_6(\sigma)$, are decreasing. The subsequences of σ , which is determined by the elements in $T_5(\sigma)$ and $T_7(\sigma)$, are increasing. So, the number of the permutations under this subcase is

$$\sum_{(T_1, T_2, T_3)} \binom{i-1}{|T_1|, |T_2|, |T_3|} 2^{|T_1|-1} 2^{|T_2|-1} 2^{|T_3|-1} 4^{j-i-1} = 2^{2j-i-6} \cdot 3(3^{i-2} - 2^{i-1} + 1),$$

where the sum is over all triples (T_1, T_2, T_3) such that $T_i \neq \emptyset$ for $i = 1, 2, 3$ and $T_1 \cup T_2 \cup T_3 = [i-1]$.

Similarly, we may consider the case $\sigma \in CP_j(\{i, j\})$ with $\sigma^{-1}(i) > \sigma^{-1}(j)$. Therefore, $cp_j(\{i, j\}) = 2[2^{j-4}(2^{i-2} - 1)(2^{j-i-1} - 1) + 2^{2j-i-6} \cdot 3(3^{i-2} - 2^{i-1} + 1)]$. In general, for any $n \geq 3$ and $3 \leq i < j \leq n$,

$$cp_n(\{i, j\}) = 2^{n-3}(2^{i-2} - 1)(2^{j-i-1} - 1) + 2^{n+j-i-5} \cdot 3(3^{i-2} - 2^{i-1} + 1).$$

■

3 The recurrence relations for the sequence $cp_n(S)$

In this section, we will derive some recurrence relations for the sequence $cp_n(S)$.

Lemma 3.1 *Let $n \geq 3$ and $S \subseteq [n-1]$. Then*

$$cp_n(S \cup \{n\}) = [n-2-2|S|]cp_{n-1}(S) + \sum_{j \notin S, j < n} 2cp_{n-1}(S \cup \{j\}).$$

Proof. Suppose $\sigma \in CP_{n-1}(S)$. We want to form a new permutation $\tau \in CP_n(S \cup \{n\})$ by inserting the letter n into σ . For any $j \in S$, since the letter j still must be a circular peak in the new permutation, we can't insert n into σ beside j . But the letter n must be a circular peak. So, there are $(n-2-2|S|)$ ways to form a new permutation τ from σ such that $\tau \in CP_n(S \cup \{n\})$.

For any $j \notin S$ with $j < n$ and $\sigma \in CP_{n-1}(S \cup \{j\})$, we must insert n into σ beside j such that n becomes a circular peak. So, there are 2 ways to form a new permutation τ from σ such that $\tau \in CP_n(S \cup \{n\})$.

Hence,

$$cp_n(S \cup \{n\}) = [n-2-2|S|]cp_{n-1}(S) + \sum_{j \notin S, j < n} 2 \cdot cp_{n-1}(S \cup \{j\}).$$

■

For any $S \in [n]$, suppose $S = \{i_1, i_2, \dots, i_k\}$, let $\mathbf{x}_S$ stand for the monomial $x_{i_1}x_{i_2} \cdots x_{i_k}$;

In particular, let $\mathbf{x}_\emptyset = 1$. Given $n \geq 3$, we define a generating function

$$g_n(x_1, x_2, \dots, x_n; y) = \sum_{\sigma \in \mathfrak{S}_n} \mathbf{x}_{CP(\sigma)} y^{|CP(\sigma)|}.$$

We also write $g_n(x_1, x_2, \dots, x_n; y)$ as g_n for short.

Corollary 3.1 *Let n be a positive integer with $n \geq 3$ and $g_n = \sum_{\sigma \in \mathfrak{S}_n} \mathbf{x}_{CP(\sigma)} y^{|CP(\sigma)|}$. Then g_n*

satisfies the following recursion:

$$g_{n+1} = [2 + (n-1)x_{n+1}y]g_n + 2x_{n+1} \sum_{i=1}^n \frac{\partial g_n}{\partial x_i} - 2x_{n+1}y^2 \frac{\partial g_n}{\partial y}.$$

for all $n \geq 3$ with initial condition $g_3 = 4 + 2x_3y$, where the notation $\frac{\partial g_n}{\partial y}$ denote partial differentiation of g_n with respect to y .

Proof. Obviously, $g_3 = 4 + 2x_3y$ and $\sum_{\sigma \in \mathfrak{S}_n} \mathbf{x}_{CP(\sigma)} y^{|CP(\sigma)|} = \sum_{S \subseteq [2, n]} cp_n(S) \mathbf{x}_S y^{|S|}$. Hence,

$$\begin{aligned}
g_{n+1} &= \sum_{S \subseteq [n+1]} cp_{n+1}(S) \mathbf{x}_S y^{|S|} \\
&= \sum_{S \subseteq [n+1], n+1 \in S} cp_{n+1}(S) \mathbf{x}_S y^{|S|} + \sum_{S \subseteq [n+1], n+1 \notin S} cp_{n+1}(S) \mathbf{x}_S y^{|S|} \\
&= \sum_{S \subseteq [n]} \left[(n-1-2|S|)cp_n(S) + \sum_{i \in [n] \setminus S} 2cp_n(S \cup \{i\}) \right] \mathbf{x}_S x_{n+1} y^{|S|+1} + 2g_n \\
&= 2 \sum_{S \subseteq [n]} \sum_{i \in [n] \setminus S} cp_n(S \cup \{i\}) \mathbf{x}_S x_{n+1} y^{|S|+1} - 2 \sum_{S \subseteq [n]} |S| cp_n(S) \mathbf{x}_S x_{n+1} y^{|S|+1} \\
&\quad + [2 + (n-1)x_{n+1}y]g_n.
\end{aligned}$$

Note that

$$\frac{\partial g_n}{\partial y} = \sum_{S \subseteq [n]} |S| cp_n(S) \mathbf{x}_S y^{|S|-1}$$

and

$$\begin{aligned}
&\sum_{S \subseteq [n]} \sum_{i \in [n] \setminus S} cp_n(S \cup \{i\}) \mathbf{x}_S x_{n+1} y^{|S|+1} \\
&= \sum_{S \subseteq [n], S \neq \emptyset} cp_n(S) x_{n+1} y^{|S|} \sum_{i \in S} \frac{\mathbf{x}_S}{x_i} \\
&= x_{n+1} \sum_{i=1}^n \frac{\partial g_n}{\partial x_i}.
\end{aligned}$$

Therefore,

$$g_{n+1} = [2 + (n-1)x_{n+1}y]g_n + 2x_{n+1} \sum_{i=1}^n \frac{\partial g_n}{\partial x_i} - 2x_{n+1}y^2 \frac{\partial g_n}{\partial y}.$$

■

Lemma 3.2 Suppose that $k \geq 0$ and $n \geq k + 4$. Let $S \subseteq [3, n - k - 1]$ with $CP_n(S) \neq \emptyset$.

Then

$$cp_n(S \cup [n - k + 1, n]) = 2(k + 1)cp_{n-1}(S \cup [n - k, n - 1]) + k(k + 1)cp_{n-2}(S \cup [n - k, n - 2]).$$

Proof. For any $\sigma \in CP_n(S \cup [n - k + 1, n])$, we consider the following four cases.

Case 1. There is no letters $i \in [n - k + 1, n]$ such that the position of i is beside $n - k$ in σ , i.e., $|\sigma^{-1}(i) - \sigma^{-1}(n - k)| = 1$. Then $\sigma^{-1}(n - k) = 1$ or n since the permutation σ hasn't a circular peak $n - k$. We can obtain a new permutation τ by exchanging the positions of $n - k$ and n in σ . Clearly, $\tau \in CP_n(S \cup [n - k, n - 1])$. Lemma 2.1 (1) tells us that $cp_n(S \cup [n - k, n - 1]) = 2cp_{n-1}(S \cup [n - k, n - 1])$. Hence, the number of permutations under this case is $2 \cdot cp_{n-1}(S \cup [n - k, n - 1])$.

Case 2. There are exact two letters $j, m \in [n - k + 1, n]$ such that $|\sigma^{-1}(j) - \sigma^{-1}(n - k)| = 1$ and $|\sigma^{-1}(m) - \sigma^{-1}(n - k)| = 1$. Deleting j and m , we obtain a subsequence τ of σ . Then $red_\sigma(\tau) \in CP_{n-2}(S \cup [n - k, n - 2])$. Note that there are $k(k - 1)$ ways to form the pairs (j, m) . Hence, the number of permutations under this case is $k(k - 1)cp_{n-2}(S \cup [n - k, n - 2])$.

Case 3. There is exact one letter $j \in [n - k + 1, n]$ such that $|\sigma^{-1}(j) - \sigma^{-1}(n - k)| = 1$. Then there are k ways to form the set $\{j\}$. Let τ be the subsequence of σ obtained by deleting j . There are the following two subcases.

Subcase 3.1. $\sigma^{-1}(n - k) \neq 1$ and n . Then $red_\sigma(\tau) \in CP_{n-1}(S \cup [n - k, n - 1])$. Hence, the number of permutations under this subcase is $k \cdot cp_{n-1}(S \cup [n - k, n - 1])$.

Subcase 3.2. $\sigma^{-1}(n - k) = 1$ or n . Then $red_\sigma(\tau) \in CP_{n-2}(S \cup [n - k, n - 2])$. Hence, the number of permutations under this subcase is $k \cdot cp_{n-2}(S \cup [n - k, n - 2])$.

So,

$$\begin{aligned}
& cp_n(S \cup [n - k + 1, n]) \\
&= 2cp_{n-1}(S \cup [n - k, n - 1]) + k(k - 1)cp_{n-2}(S \cup [n - k, n - 2]) \\
&\quad + 2k \cdot cp_{n-1}(S \cup [n - k, n - 1]) + 2k \cdot cp_{n-2}(S \cup [n - k, n - 2]) \\
&= 2(k + 1)cp_{n-1}(S \cup [n - k, n - 1]) + k(k + 1)cp_{n-2}(S \cup [n - k, n - 2]).
\end{aligned}$$

■

Setting $S = \emptyset$ in Lemma 3.2, we derive the following results.

Theorem 3.1 *Let $k \geq 1$, $n \geq \max\{3, 2k\}$ and $CP_n([n - k + 1, n]) \neq \emptyset$. Suppose that*

$$cp_n([n - k + 1, n]) = k(k + 1) \sum_{i=1}^k (-1)^{i+1} b_{k,i} [(2k + 2)^{n-2k} - (2k + 2 - 2i)^{n-2k}].$$

Then the coefficients $b_{k,i}$ satisfy the recurrence relation as follows:

$$b_{k+1,i} = \begin{cases} k(k + 1)^2 \sum_{j=1}^k (-1)^{j+1} b_{k,j} & \text{if } i = 1 \\ k(k + 1) \frac{k+2-i}{i} b_{k,i-1} & \text{if } 2 \leq i \leq k + 1 \end{cases}.$$

Proof. Lemma 2.1 tells us that $cp_n([n, n]) = cp_n(\{n\}) = 4^{n-2} - 2^{n-2}$. Hence, $b_{1,1} = \frac{1}{2}$.

Lemma 3.2 implies that

$$cp_n([n - k + 1, n]) = 2(k + 1)cp_{n-1}([n - k, n - 1]) + k(k + 1)cp_{n-2}([n - k, n - 2]).$$

By comparing the coefficients, we obtain the desired results. ■

By Theorem 3.1, we compute the values of $b_{k,i}$ for all $1 \leq k \leq 4$ as follows.

	$i = 1$	2	3	4
$k = 1$	$\frac{1}{2}$			
2	2	$\frac{1}{2}$		
3	27	12	1	
4	768	486	96	3

Table.1. The values of $b_{k,i}$ for $1 \leq k \leq 4$

Corollary 3.2 *Let $k \geq 0$, $n \geq \max\{3, 2k\}$ and $CP_n([n - k + 1, n]) \neq \emptyset$. Suppose that*

$$cp_n([n - k + 1, n]) = \sum_{i=0}^k (-1)^i a_{k,i} (2k + 2 - 2i)^{n-2k}.$$

Then the coefficients $a_{k,i}$ satisfy the following recurrence relation:

$$a_{k+1,i} = \begin{cases} \sum_{j=1}^{k+1} (-1)^{j+1} a_{k+1,j} & \text{if } i = 0 \\ (k+1)(k+2) \frac{k+2-i}{i} a_{k,i-1} & \text{if } 1 \leq i \leq k+1 \end{cases},$$

with initial condition $a_{0,0} = \frac{1}{2}$. Let $f_k(x) = \sum_{i=0}^k a_{k,i} x^i$ for any $k \geq 0$, then $f_0(x) = \frac{1}{2}$, $f_{k+1}(-1) = 0$ and $f_k(x)$ satisfies the differential equation $f'_{k+1}(x) = (k+1)^2(k+2)f_k(x) - (k+1)(k+2)xf'_k(x)$ for any $k \geq 0$, where the notation “'” denotes the differentiation of functions.

Proof. When $k = 0$, by Lemma 2.1, we have $cp_n([n + 1, n]) = cp_n(\emptyset) = 2^{n-1}$. Hence, $a_{0,0} = \frac{1}{2}$. When $k \geq 1$, it is easy to check that

$$a_{k,i} = \begin{cases} k(k+1) \sum_{j=1}^k (-1)^{j+1} b_{k,j} & \text{if } i = 0 \\ k(k+1)b_{k,i} & \text{if } 1 \leq i \leq k \end{cases}.$$

Hence, by Theorem 3.1, we have

$$a_{k+1,i} = \begin{cases} \sum_{j=1}^{k+1} (-1)^{j+1} a_{k+1,j} & \text{if } i = 0 \\ (k+1)(k+2) \frac{k+2-i}{i} a_{k,i-1} & \text{if } 1 \leq i \leq k+1 \end{cases},$$

with initial condition $a_{0,0} = \frac{1}{2}$. Let $f_k(x) = \sum_{i=0}^k a_{k,i} x^i$ for any $k \geq 0$. Clearly, $f_0(x) = \frac{1}{2}$.

Note that $a_{k,i} = \sum_{j=1}^k (-1)^{j+1} a_{k,j}$ for any $k \geq 1$. Hence, $f_{k+1}(-1) = 0$ if $k \geq 0$. Since $a_{k+1,i} = (k+1)(k+2) \frac{k+2-i}{i} a_{k,i-1}$ for any $i \in [k+1]$, we have

$$\sum_{i=1}^{k+1} a_{k+1,i} x^i = \sum_{i=1}^{k+1} (k+1)(k+2) \frac{k+2-i}{i} a_{k,i-1} x^i$$

Simple computations tell us that $f'_{k+1}(x) = (k+1)^2(k+2)f_k(x) - (k+1)(k+2)xf'_k(x)$. ■

By Corollary 3.2, we compute the values of $a_{k,i}$ for all $0 \leq k \leq 3$ as follows.

	$i = 0$	1	2	3
$k = 0$	$\frac{1}{2}$			
1	1	1		
2	9	12	3	
3	192	324	144	12

Table.2. The values of $a_{k,i}$ for $0 \leq i \leq 3$

Recall that $w(i, r, n, k) = \sum_{P \in P_{r,n,k}} w_i(P)$, where $P_{r,n,k}$ is the set of all the circular-peak path from the vertices $(r, 0)$ to (n, k) and $w_i(P)$ is the weight of path $P \in P_{r,n,k}$.

Lemma 3.3

$$w(i, r, n, k) = 2^{n-r-2k} \prod_{m=0}^{k-1} (m+i+1)(m+i+2) \sum \prod_{m=0}^k (i+m+1)^{t_m}$$

where the sum is over all $(k+1)$ -tuples $(t_0, t_1, \dots, t_k)$ such that $\sum_{m=0}^k t_m = n - r - 2k$ and $t_m \geq 0$.

Proof. Suppose that $P = e_1 e_2 \cdots e_{n-k-r} \in P_{r,n,k}$. Let $\mathcal{R} = \{j \mid e_j = R\}$, then $|\mathcal{R}| = k$. Furthermore, we may suppose that $\mathcal{R} = \{e_{j_1}, \dots, e_{j_k}\}$, where $0 = j_0 < j_1 < j_2 < \dots < j_k \leq n - k - r = j_{k+1}$. Hence,

$$w_i(P) = \prod_{m=0}^k [2i + 2m + 2]^{j_{m+1} - j_m - 1} \prod_{m=0}^{k-1} (m + i + 1)(m + i + 2).$$

Let $t_m = j_{m+1} - j_m - 1$ for any $0 \leq m \leq k$, then $t_m \geq 0$ and $\sum_{m=0}^k t_m = n - r - 2k$. So,

$$\begin{aligned} w(i, r, n, k) &= \sum \prod_{m=0}^k [2i + 2m + 2]^{t_m} \prod_{m=0}^{k-1} (m + i + 1)(m + i + 2) \\ &= 2^{n-r-2k} \prod_{m=0}^{k-1} (m + i + 1)(m + i + 2) \sum \prod_{m=0}^k [i + m + 1]^{t_m} \end{aligned}$$

where the sum is over all $(k+1)$ -tuples $(t_0, t_1, \dots, t_k)$ such that $\sum_{m=0}^k t_m = n - r - 2k$ and $t_m \geq 0$. ■

Lemma 3.4 Suppose $n \geq 3$ and $k \geq 0$. Let $S \subseteq [3, n - k - 1]$ with $CP_n(S) \neq \emptyset$ and $r = \max S$. Then

$$cp_n(S \cup [n - k + 1, n]) = \sum_{i=0}^k w(i, r, n - i, k - i) cp_{r+i}(S \cup [r + 1, r + i]).$$

Proof. We view the set $S \cup [n - k + 1, n]$ as a vertex (n, k) . Connect the vertices (n, k) with $(n - 1, k)$ (*resp.* $(n - 2, k - 1)$) and give this edge a weight $2(k + 1)$ (*resp.* $k(k + 1)$). We draw the obtained graph as follows:





Fig.1. the obtained graph from the recurrence relation in Lemma 3.2

Conversely, from the graph, we also can derive the recurrence relation in Lemma 3.2. Repeating this processes, we will obtain the following graph with weight.



Hence, $cp_n(S \cup [n - k + 1, n])$ can be expressed as a linear combination of $cp_{r+i}(S \cup [r + 1, r])$, $cp_{r+i}(S \cup [r + 1, r + 1])$, $\dots$, and $cp_{r+i}(S \cup [r + 1, r + k])$. It is easy to check that the coefficient of $cp_{r+i}(S \cup [r + 1, r + i])$ is $w(i, r, n - i, k - i)$ which is the sum of the weights of all the circular peak path from $(r + i, i)$ to (n, k) . So, $cp_n(S \cup [n - k + 1, n]) = \sum_{i=0}^k w(i, r, n - i, k - i) cp_{r+i}(S \cup [r + 1, r + i])$. ■

For any $S \subseteq [3, n]$, suppose that the type of the set S is $(r_1^{k_1}, r_2^{k_2}, \dots, r_m^{k_m})$. We have studied the case with $m = 1$ in Theorem 3.1. We will consider the case with $m \geq 2$ in the following theorem.

Theorem 3.2 *Let $n \geq 3$, $k \geq 0$ and $S \subseteq [3, n]$. Suppose that $CP_n(S) \neq \emptyset$ and the type of*

the set S is $(r_1^{k_1}, r_2^{k_2}, \dots, r_m^{k_m})$ with $m \geq 2$, where $m \geq 2$. Then

$$cp_n(S) = 2^{n-r_m} \sum_{i_1=0}^{k_m} \sum_{i_2=0}^{k_{m-1}+i_1} \cdots \sum_{i_{m-1}=0}^{k_2+i_{m-2}} \prod_{j=1}^{m-1} w(i_j, r_{m-j}, r_{m-j+1} + i_{j-1} - i_j, k_{m-j+1} + i_{j-1} - i_j) cp_{r_1+i_{m-1}}([r_1 - k_1 + 1, r_1 + i_{m-1}])$$

where $i_0 = 0$.

Proof. Iterating the identity in Lemma 3.4, we obtain the desired results. ■

Example 3.1 Let $S = \{i, j\}$ such that $3 \leq i < j - 1$, $j \leq n$ and $CP_n(S) \neq \emptyset$. The type of S is (i^1, j^1) . By Theorem 3.2, we have $cp_n(\{i, j\}) = 2^{n-j}[w(0, i, j, 1)cp_i(\{i\}) + w(1, i, j - 1, 0)cp_{i+1}(\{i, i + 1\})]$. Lemma 3.3 implies that $w(0, i, j, 1) = 2^{j-i-1}(2^{j-i-1} - 1)$ and $w(1, i, j - 1, 0) = 2^{2j-2i-2}$. Theorem 2.1 tells us that $cp_i(\{i\}) = 2^{i-1}(2^{i-1} - 1)$. From Corollary 3.2 it follows that $cp_{i+1}(\{i, i + 1\}) = \sum_{m=0}^2 (-1)^m a_{2,m} (6 - 2m)^{i-3} = 9 \cdot 6^{i-3} - 12 \cdot 4^{i-3} + 3 \cdot 2^{i-3}$. Hence, we obtain the formula $cp_n(\{i, j\}) = 2^{n-3}(2^{i-2} - 1)(2^{j-i-1} - 1) + 2^{n+j-i-5} \cdot 3(3^{i-2} - 2^{i-1} + 1)$ again.

4 Appendix

For convenience to check the identities given in the previous sections, by the computer search, for $3 \leq n \leq 8$, we obtain the number $cp_n(S)$ of the permutations in the sets $CP_n(S) \neq \emptyset$ and list them in Table 3.

$n = 3$	$S = \emptyset$	$\{3\}$							
	4	2							
$n = 4$	$S = \emptyset$	$\{3\}$	$\{4\}$						
	8	4	12						
$n = 5$	$S = \emptyset$	$\{3\}$	$\{4\}$	$\{5\}$	$\{3, 5\}$	$\{4, 5\}$			
	16	8	24	56	4	12			
$n = 6$	$S = \emptyset$	$\{3\}$	$\{4\}$	$\{5\}$	$\{6\}$	$\{3, 5\}$	$\{3, 6\}$	$\{4, 5\}$	$\{4, 6\}$
	32	16	48	112	240	8	24	24	72
	$S = \{5, 6\}$								
	144								
$n = 7$	$S = \emptyset$	$\{3\}$	$\{4\}$	$\{5\}$	$\{6\}$	$\{7\}$	$\{3, 5\}$	$\{3, 6\}$	$\{3, 7\}$
	64	32	96	224	480	992	16	48	112
	$S = \{4, 5\}$	$\{4, 6\}$	$\{4, 7\}$	$\{5, 6\}$	$\{5, 7\}$	$\{6, 7\}$	$\{3, 5, 7\}$	$\{3, 6, 7\}$	$\{4, 5, 7\}$
	48	144	336	288	688	1200	8	24	24
	$S = \{4, 6, 7\}$	$\{5, 6, 7\}$							
	72	144							
$n = 8$	$S = \emptyset$	$\{3\}$	$\{4\}$	$\{5\}$	$\{6\}$	$\{7\}$	$\{8\}$	$\{3, 5\}$	$\{3, 6\}$
	128	64	192	448	960	1984	4032	32	96
	$S = \{3, 7\}$	$\{3, 8\}$	$\{4, 5\}$	$\{4, 6\}$	$\{4, 7\}$	$\{4, 8\}$	$\{5, 6\}$	$\{5, 7\}$	$\{5, 8\}$
	224	480	96	288	672	1440	576	1376	2976
	$S = \{6, 7\}$	$\{6, 8\}$	$\{7, 8\}$	$\{3, 5, 7\}$	$\{3, 5, 8\}$	$\{3, 6, 7\}$	$\{3, 6, 8\}$	$\{3, 7, 8\}$	$\{4, 5, 7\}$
	2400	5280	8640	16	48	48	144	288	48
	$S = \{4, 5, 8\}$	$\{4, 6, 7\}$	$\{4, 6, 8\}$	$\{4, 7, 8\}$	$\{5, 6, 7\}$	$\{5, 6, 8\}$	$\{5, 7, 8\}$	$\{6, 7, 8\}$	
	144	144	432	864	288	864	1728	2880	

Table.3. $cp_n(S)$ for $3 \leq n \leq 8$ with $CP_n(S) \neq \emptyset$

References

- [1] Pierre Bouchard, Jun Ma, Yeong-Nan Yeh, Circular Peaks and Hilbert Series,
- [2] Hungyung Chang, Jun Ma, Yeong-Nan Yeh, Enumerations of Permutations by Circular Descent Sets
- [3] Robert J. Clarke, Einar Steingrímsson, Jiang Zeng, New Euler-Mahonian permutation statistics, *Sm. Lothar. Combin.* 35 (1995), Art. B35c, approx. 29 pp. (electronic).

- [4] Michael Domaratzki, Combinatorial Interpretations of a Generalization of the Genocchi Numbers, *Journal of Integer Sequences*, Vol. 7 (2004), Article 04.3.6
- [5] J. R. Stembridge, Enriched P-partitions, *Trans. Amer. Math. Soc.* 349 (1997), no. 2, 763-788.